

Proof of your article (# A1135-4) from "Journal of Applied Physiology" is available for download

=====

Dear Sir or Madam:

Please refer to this URL address

<http://rapidproof.cadmus.com/RapidProof/retrieval/index.jsp>

Login: your e-mail address

Password: ----

The file at the above URL address contains the following:

- Proofreading marks
- Reprint Order form
- Copyedited page proof of your article

The site contains 1 file. You will need Adobe Acrobat® Reader to read this file. Adobe Acrobat® Reader is a free software, available for user downloading at <http://www.adobe.com/products/acrobat/readstep.html>.

After you print the PDF file of your paper, please read the page proof carefully and

- 1) clearly indicate all changes or corrections on the margin;
- 2) answer all queries (footnotes 1, 2, 3, etc.) listed on the last page of the PDF proof;
- 3) carefully proofread all/any tables and equations;
- 4) make sure that any special characters, such as Greek letters, especially μ (mu), have translated correctly;
- 5) If you have questions about figure quality, note your concerns on the margin of the relevant page. Please keep in mind that the final printed version will be of higher quality than the PDF proof and that the online version of the published article will appear identical.

IMPORTANT: Do NOT make online alterations to the PDF itself. Corrections within the PDF, made with the Adobe Acrobat-provided tools will remain unseen by the Journal staff. This will result in the final article NOT reflecting your intended changes. ONLY the hand-marked PDF printout guarantees that the changes will be incorporated into the final text of the article.

We encourage you to retain a copy of the proof with your corrections, should further changes and/or clarifications be required.

IMPORTANT NOTES

1. To guarantee the placement of your article in the next available issue of the "Journal of Applied Physiology", please return the corrected set of PDF page proof via an overnight courier service to this address WITHIN 48 HOURS:

The American Physiological Society
"Journal of Applied Physiology" PROOF
9650 Rockville Pike
Bethesda, MD 20814-3991
USA
phone: 301-634-7070

2. If you require a hard copy of your color image(s), you may request it by replying to this message. The color proof will then be provided to you at an additional fee of \$75.00 per color figure, which will be added to the publication fees for your article. You must provide the appropriate figure number(s) and your mailing address in your reply if you wish to receive the hard copy of the color proof(s).

3) The filled out Reprint Form must be returned within 24 hours to the address below:

Cadmus Professional Communications
Cadmus Reprints
PO Box 751903
Charlotte NC 28275-1903
USA

For reprint inquiries, please contact Mary Leonard, fax: 410-770-4659, e-mail: leonardm@cadmus.com.

If you have any problems or questions, please contact me.

PLEASE ALWAYS INCLUDE YOUR ARTICLE NO. (A1135-4) WITH ALL CORRESPONDENCE.

Sincerely,

Virginia Million
Journal Editorial Supervisor
Journal of Applied Physiology

9650 Rockville Pike
Bethesda, MD 20814
Tel: 301-634-7216
Fax: 301-634-7243
E-mail: vmillion@the-aps.org

Proofreader's Marks

MARK	EXPLANATION	EXAMPLE
	TAKE OUT CHARACTER INDICATED	Your proof.
^	LEFT OUT, INSERT	u Yor proof. ^
#	INSERT SPACE	# Yourproof. ^
9	TURN INVERTED LETTER	Your p ^l oof. ^
X	BROKEN LETTER	X Your pr ^o of.
eg#	EVEN SPACE	eg# A good proof.
C	CLOSE UP: NO SPACE	Your pro ^o gf.
tr	TRANSPOSE	tr A proof good
wf	WRONG FONT	wf Your proof.
lc	LOWER CASE	lc Your proof.
≡ caps	CAPITALS	Your proof. caps Your proof.
ital	ITALIC	Your proof. ital Your proof.
rom	ROMAN, NON ITALIC	rom Your proof.
bf	BOLD FACE	Your proof. bf Your proof.
..... stet	LET IT STAND	Your proof. stet Your proof.
out sc.	DELETE, SEE COPY	out sc. She Our proof. ^
spell out	SPELL OUT	spell out Queen (Eliz.)
#	START PARAGRAPH	# read. [Your
no #	NO PARAGRAPH: RUN IN	no # marked. → # Your proof.
L	LOWER	L [Your proof.]

MARK	EXPLANATION	EXAMPLE
┌	RAISE	┌ Your proof.
└	MOVE LEFT	└ Your proof.
┐	MOVE RIGHT	┐ Your proof.
	ALIGN TYPE	└ Three dogs. Two horses.
≡	STRAIGHTEN LINE	≡ Your proof.
⊙	INSERT PERIOD	⊙ Your proof. ^
;/	INSERT COMMA	;/ Your proof. ^
:/	INSERT COLON	:/ Your proof. ^
;/	INSERT SEMICOLON	;/ Your proof. ^
∨	INSERT APOSTROPHE	∨ Your mans proof. ^
∨ ∨	INSERT QUOTATION MARKS	∨ ∨ Marked it proof. ^ ^
=/	INSERT HYPHEN	=/ A proofmark. ^
!	INSERT EXCLAMATION MARK	! Prove it. ^
?	INSERT QUESTION MARK	? Is it right. ^
Ⓚ	QUERY FOR AUTHOR	Ⓚ was Your proof read by ^
[/]	INSERT BRACKETS	[/] The Smith girl ^ ^
</>	INSERT PARENTHESES	</> Your proof. ^ ^
1/m	INSERT 1-EM DASH	1/m Your proof. ^
□	INDENT 1 EM	□ Your proof
▢	INDENT 2 EMS	▢ Your proof.
▣	INDENT 3 EMS	▣ Your proof.

Journal of Applied Physiology 2005

Published by The American Physiological Society

This is your pro forma invoice or order form

(Please keep a copy of this document for your records. This form is not for commercial ordering)

IMPORTANT Order form must be returned within 48 hours of receipt to avoid late charges. Orders received after 48 hours will be charged an additional fee of 25%. Orders received after 30 days will be charged an additional 50%. Reprints containing color figures are available only if ordered before the journal is printed. It is the policy of Cadmus Reprints to issue only one invoice per order. Please print clearly.

Author Name _____
Title of Article _____
Issue of Journal _____ Manuscript # A1135-4 Publication Date _____
Number of Pages _____ Article # _____ Symbol JAP
Color in Article? Yes / No (Please Circle)

Please include the journal name and reprint number or manuscript number on your purchase order or other correspondence.

Reprint and Publication Costs

Reprint Costs (Please see page 2 for reprint costs/fees.)

Number of reprints ordered _____ \$ _____
Add color in reprints: \$100 per
100 copies up to 500 copies \$ _____
Subtotal \$ _____
Add appropriate sales tax/GST to subtotal \$ _____
Add \$32 for each additional ship location \$ _____
Total Reprint Due \$ _____

Publication Fees (Please see page 2 for reprint costs/fees.)

Page Charges: \$70 per journal page \$ _____
Color Figures: \$300 per color figure \$ _____
Hard copy color proof: \$75 per figure \$ _____
Toll-Free Link: \$150 \$ _____

Member No. _____ Member Signature _____
Total Publication Fees \$ _____
TOTAL TO REMIT \$ _____

Payment Details (FEIN #:541274108)

Enclosed: Personal Check _____
Institutional Purchase Order _____
Credit Card Payment Details _____
Checks must be paid in U.S. dollars and drawn on a U.S. Bank.
Credit Card: ___ VISA ___ Am. Exp. ___ MasterCard
Card Number _____
Expiration Date _____
Signature _____

Cadmus will process credit cards and Cadmus Journal Services will appear on the credit card statement.

Send payment and order form (no express packages) to:

Cadmus Reprints Fax order form
P.O. Box 751903 to 410-820-9765
Charlotte, NC 28275-1903

Invoice and Shipping Information

Invoice Address

Name _____
Institution _____
Department _____
Street _____
City _____ State _____ Zip _____
Country _____
Phone _____ Fax _____
E-mail Address _____
Purchase Order No. _____

Shipping Address (cannot ship to a P.O. Box.)

Name _____
Institution _____
Department _____
Street _____
City _____ State _____ Zip _____
Country _____
Quantity _____ Fax _____
Phone: Day _____ Evening _____

Additional Shipping Address* (cannot ship to a P.O. Box)

Name _____
Institution _____
Department _____
Street _____
City _____ State _____ Zip _____
Country _____
Quantity _____ Fax _____
Phone: Day _____ Evening _____

* Add \$32 for each additional shipping address

SIGNATURE REQUIRED: By signing this form the author agrees to accept responsibility for the payment of the mandatory page charges of \$70 per page, reprints ordered, as well as any color charges, late payments, and split shipment charges. If the charges are billed to an institution, the author must assume the responsibility for making the necessary arrangements for the issuance of a formal institutional purchase order. Otherwise, it is understood that the author will bear the cost of these charges. Failure to pay any of these agreed-upon charges could jeopardize future submissions.

AUTHOR Signature _____
Telephone _____

Fax _____
E-mail _____

Journal of Applied Physiology (JAP)

Published by The American Physiological Society
2005 REPRINT AND PUBLICATION CHARGES
Author rates only.
Not to be used for commercial ordering.

Black and White Reprint Prices

Domestic (USA only)						
# of Pages	100	200	300	400	500	Addl 100's
1-4	\$196	\$273	\$350	\$427	\$504	\$71
5-8	\$265	\$401	\$539	\$675	\$811	\$122
9-12	\$342	\$520	\$702	\$879	\$1,059	\$137
13-16	\$410	\$649	\$887	\$1,126	\$1,366	\$227
17-20	\$477	\$769	\$1,059	\$1,351	\$1,638	\$278
21-24	\$555	\$897	\$1,237	\$1,579	\$1,919	\$325
25-28	\$622	\$1,025	\$1,426	\$1,827	\$2,227	\$382
29-32	\$702	\$1,152	\$1,615	\$2,074	\$2,535	\$438

International (includes Canada and Mexico)						
# of Pages	100	200	300	400	500	Addl 100's
1-4	\$222	\$310	\$401	\$490	\$582	\$83
5-8	\$302	\$464	\$627	\$790	\$951	\$148
9-12	\$393	\$608	\$833	\$1,045	\$1,267	\$175
13-16	\$474	\$763	\$1,053	\$1,345	\$1,637	\$279
17-20	\$554	\$910	\$1,267	\$1,621	\$1,972	\$342
21-24	\$643	\$1,063	\$1,483	\$1,902	\$2,321	\$403
25-28	\$726	\$1,217	\$1,707	\$2,202	\$2,690	\$473
29-32	\$817	\$1,372	\$1,938	\$2,502	\$3,067	\$542

Minimum order is 100 copies. For articles longer than 32 pages, please consult Cadmus Reprints at 800-407-9190.

Page Charges

Add \$70 per journal page for all pages in the article.

Color

Reprints containing color figures are available only if ordered before journal is printed. Add \$100 per 100 copies to the cost of the reprints if the article contains color.

If your article contains **color**, you must pay subsidized color charges of \$300/fig. (reprint charge is \$1000/fig for those who do not pay promptly), whether or not you buy reprints. These **color charges are waived for APS Members who are the first or last author of the paper.**

If you requested a **hard copy color figure proof** when you reviewed your S-proof, the charge is \$75.

TOLL-FREE LINK

A link can be created from a url of your choice to your article online so that readers accessing your article from your url can do so without a subscription. The cost is \$150. This is especially useful if your article contains electronic supplemental material.

For more information, please click on this link:
<http://www.the-aps.org/publications/sprooflink.pdf>

Shipping

Shipping costs are included in the reprint prices. Domestic orders are shipped via UPS Ground service. Foreign orders are shipped via an expedited air service. The shipping address printed on an institutional purchase order always supercedes.

Multiple Shipments

Orders can be shipped to more than one location. Please be aware that it will cost \$32 for each additional location.

State Sales Tax and Canadian GST

Residents of Virginia, Maryland, Pennsylvania, and the District of Columbia are required to add the appropriate sales tax to each reprint order. For orders shipped to Canada, please add 7% Canadian GST unless exemption is claimed.

Ordering

Please fax your order form and purchase order to 410-820-9765. Prepayment of checks should be mailed to address below:

Cadmus Reprints
P.O. Box 751903
Charlotte, NC 28275-1903

*Note: Do not send express packages to this location.
FEIN #:541274108*

Please direct all inquiries to:

Mary Leonard
800-407-9190 (toll free number)
410-819-3912 (direct number)
410-820-9765 (FAX number)
LeonardM@cadmus.com

HIGHLIGHTED TOPIC | *Biomechanics and Mechanotransduction in Cells and Tissues***High-frequency, low-magnitude vibrations suppress the number of blood vessels per muscle fiber in mouse soleus muscle**

AQ:1

Walter L. Murfee,¹ Laura A. Hammett,¹ Caroline Evans,¹ Liqin Xie,² Maria Squire,² Clinton Rubin,² Stefan Judex,² and Thomas C. Skalak¹¹Department of Biomedical Engineering, University of Virginia Health Sciences Center, Charlottesville, Virginia; and ²Department of Biomedical Engineering, State University of New York at Stony Brook, Stony Brook, New York

Submitted 8 October 2004; accepted in final form 18 January 2005

Murfee, Walter L., Laura A. Hammett, Caroline Evans, Liqin Xie, Maria Squire, Clinton Rubin, Stefan Judex, and Thomas C. Skalak. High-frequency, low-magnitude vibrations suppress the number of blood vessels per muscle fiber in mouse soleus muscle. *J Appl Physiol* 98: 000–000, 2005. First published January 27, 2005; doi:10.1152/jappphysiol.01135.2004.—Extremely low-magnitude (0.3 g), high-frequency (30–90 Hz), whole body vibrations can stimulate bone formation and are hypothesized to provide a surrogate for the oscillations of muscle during contraction. Little is known, however, about the potential of these mechanical signals to stimulate adaptive responses in other tissues. The objective of this study was to determine whether low-level mechanical signals produce structural adaptations in the vasculature of skeletal muscle. Eight-week-old male BALB/cByJ (BALB) mice were divided into two experimental groups: mice subjected to low-level, whole body vibrations (45 Hz, 0.3 g) superimposed on normal cage activities for 15 min/day ($n = 6$), and age-matched controls ($n = 7$). After the 6-wk experimental protocol, sections from end and mid regions of the soleus muscles were stained with BSI lectin, an endothelial cell marker, and smooth muscle (SM) α -actin, a perivascular cell marker. Six weeks of this low-level vibration caused a 29% decrease in the number of lectin-positive vessels per muscle fiber in the end region of the soleus muscle, indicating a significant reduction in the number of capillaries per muscle fibers. Similarly, these vibrations caused a 36% reduction in SM α -actin-positive vessels per muscle fiber, indicating a reduction in the number of arterioles and venules. The decreases in lectin- and SM α -actin-positive vessels per muscle fiber ratios were not significant in the mid muscle sections. These results demonstrate the sensitivity of the vasculature in mouse skeletal muscle to whole body, low-level mechanical signals.

AQ:2

AQ:3 microcirculation; angiogenesis; musculoskeletal adaptation

MECHANICAL SIGNALS ARE CRITICAL to the regulation of a wide spectra of vertebrate systems, including gene expression, cell growth, tissue architecture, organ morphology and even function of the organism. However, the critical frequency and amplitude ranges for influencing the plasticity of the response remains elusive. Recent evidence has indicated that extremely low-level mechanical signals, delivered to the weight-bearing skeleton in the form of vibration, are anabolic to bone (30, 31). This provides evidence that bone tissue, often thought to be anabolically responsive only to those strains generated during extreme physical activity (2,000–3,000 microstrain), may in

fact be sensitive to the extremely low-level, high-frequency mechanical signals that arise through normal daily activities. More specifically, 30- to 90-Hz vibrations applied for as little as 10 min/day at one-third of the Earth's acceleration (0.3 g) induce bone strains two orders of magnitude less than peak strains experienced during normal locomotion, yet still lead to increased bone formation, reduced bone loss, and enhanced mechanical properties in various species including sheep, mice, rats, and humans (17, 29–33, 39). Similarly, whole body vibration can be used to improve muscular performance, balance, and neurosensing (6, 27, 34, 36, 38). Despite the clinical potential of low-magnitude, whole body vibration to provide a nonpharmacological intervention for diseases such as osteoporosis, the impact of these mechanical signals on the metabolism and morphology of other tissues remains unknown.

Muscle is also known to experience relatively high-frequency mechanical stimuli. During muscle fiber contraction, there is strong evidence of mechanical energy in the frequency domain of 10–100 Hz. This same frequency domain is also evident in the functional strain history of the skeleton (10), with a considerable portion of the mechanical regulatory information that a bone is subject to derived from these persistent mechanical signals. In addition to bone's sensitivity to mechanical signals in this range, muscle spindle primary endings have been shown to be sensitive to high-frequency vibrations (3, 4, 22). High-frequency, low-amplitude (0.3 mm) tendon vibration for less than 4 min/day can attenuate hindlimb unloading effects in rat soleus muscle (8).

The effects of mechanical stimuli on the musculature can also produce changes in the vasculature. Externally applied mechanical forces elicited by stretching, electrical stimulation of muscles, or internally generated forces by exercise are associated with vascular adaptations at the arteriole and capillary level (1, 2, 7, 13, 26). Depending on the type and magnitude of the stimulus, hemodynamic alterations can cause vascular cell remodeling, including differentiation and proliferation, leading to network growth and in some cases rarefaction (2, 13, 23, 24, 25, 37), and indicate the ability of adult microvasculature to adapt to imposed epigenetic strains of magnitudes and frequencies in the range of 1–20% strain and 1–10 Hz (5, 19, 35). However, the ability of the microvasculature to respond to mechanical signals outside these ranges is

AQ:8 Address for reprint requests and other correspondence: W. L. Murfee, Dept. of Biomedical Engineering, Box 800759, UVA Health System, Charlottesville, VA 22908 (E-mail: wlm5e@virginia.edu).

AQ:8 The costs of publication of this article were defrayed in part by the payment of page charges. The article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

essentially unknown. Recognizing that whole body vibrations are transduced from the skeleton to attached postural muscles and muscle is able to sense vibration, we postulated that whole body vibration could have an effect on the vasculature in muscle. Such an effect might be mediated via direct physical transduction of muscle strains to the vascular cells or via an anabolic/phenotypic adaptation of the muscle that leads to alterations in blood supply.

On the basis of the use of the number of blood vessels per muscle fiber as a metric of microvascular remodeling in skeletal muscle in response to mechanical stimulation (1, 2, 7, 13), the objective of this study was to assess the effect of high-frequency, low-magnitude vibrations on the number of capillaries and arterioles per muscle fiber. BSI lectin was used to identify endothelial cells lining microvessels, whereas an antibody against smooth muscle (SM) α -actin was used to label perivascular cells, defined as vascular mural cells wrapping around or elongated along the endothelium, along arterioles and venules. We hypothesized that high-frequency, low-magnitude, whole body vibrations, known to be anabolic to bone, could increase the number of vessels per muscle fiber in skeletal muscle. Our findings for the first time suggest that low-level, whole body vibrations are capable of affecting the microvasculature within skeletal muscle.

METHODS

Experimental protocol. All experimental procedures were performed in accordance with the guidelines of the State University of New York at Stony Brook Animal Care and Use Committee. Representing a model for the young adult musculoskeletal system, eight wk-old male BALB/cByJ (BALB) mice (Jackson Laboratory, Bar Harbor, ME) were divided into two experimental groups: age-matched control (AC; $n = 10$) and mechanically stimulated (MS; $n = 10$). Inbred BALB mice were chosen for this study because of sensitivity of their skeleton to whole body vibration (17, 18). Each mouse was housed in an individual plastic cage. For mice in the MS group, whole body vibrations were applied by placing the individual plastic cages on a plate, which was vertically oscillating in a sinusoidal fashion at 45 Hz, producing peak accelerations of 0.3 g (Refs. 9, 17; 1.0 g = Earth's gravitational field or 9.8 ms⁻²). Vertical peak-to-peak displacement of the cage amounted to 74 μ m, making the displacement barely perceptible to human touch. Each mouse was allowed to freely move within its cage at all times, including during the 15-min vibration session, thus superimposing the vibratory external mechanical signal onto normal cage activity. After the 6-wk protocol, mice were euthanized by carbon dioxide inhalation, and the right soleus muscle was harvested from each animal, sectioned, and stained for BSI lectin and SM α -actin.

Tissue harvesting and sectioning. The soleus muscle from the right limb was dissected from the posterior surface of the tibia and pinned to approximately retain the original *in vivo* muscle length. The pinned muscle specimens were embedded in TBS tissue freezing medium (Triangle Biomedical Sciences), frozen in dry ice-cooled isopentane, and stored in an isopentane-filled Eppendorf tube at -80°C until sectioning. Before being sectioned, each specimen was embedded and re-snap frozen in Tissue-Tek OCT. Each soleus muscle was oriented to ensure that sectioning would begin at the distal end. Multiple frozen sections were cut from the end (1,000 μ m into the muscle) and mid region (4,500 μ m into the muscle) of each muscle specimen. Sections were mounted onto gelatin-coated slides and fixed in 100% methanol for 30 min at -20°C.

Immunocytochemistry. The number of lectin-positive and SM α -actin-positive vessels per muscle fiber, the number of muscle fibers per area, and the total muscle cross-sectional area were evaluated in both the

end and mid-region sections of the muscle. Acceptable specimens were determined by satisfaction of two criteria: 1) positive vascular staining of BSI lectin and SM α -actin across the entire cross section of the muscle and 2) muscle sections were obtained from both the end and mid regions of the muscle. Samples that failed to satisfy these two criteria were not included for analysis. Consequently, three samples from the control group and four samples from the mechanically stimulated group were excluded, resulting in $n = 7$ for control (AC) and $n = 6$ for MS. BSI lectin was used to identify endothelial cells lining microvessels, whereas an antibody against SM α -actin was used to label perivascular cells, defined as vascular mural cells wrapping around or elongated along the endothelium. This includes pericytes and mature and immature SM cells. BSI lectin was previously characterized as a marker of capillaries in sectioned muscle (14). In our experience, BSI lectin labels all capillaries including angiogenic sprouts in skeletal muscle and rat mesentery. In contrast, SM α -actin labels SM cells wrapping around arterioles and venules and is also expressed by a population of pericytes found elongated along capillaries. SM α -actin is not expressed by pericytes along every capillary and thus has been used as a marker of arterioles, venules, and capillaries with differentiated pericytes (13, 37). Additionally, end soleus muscle sections from control and mechanically stimulated mice were assessed for myocyte myosin ATPase activity (pH 4.3) to determine muscle fiber type ($n = 3$ for AC group, $n = 4$ for MS group).

After tissue harvesting and sectioning, muscle sections were washed in PBS for 30 min and incubated in 1:20 BSI lectin-FITC (Sigma), 1:200 IA4-CY3 (Sigma) in antibody buffer (PBS with 0.1% saponin, 2% bovine serum albumin) for 1 h at room temperature. Sections were then washed every 15 min for 30 min in PBS containing 0.1% saponin. After tissue staining, sections were examined with a Nikon TE 300 microscope equipped with confocal accessories (Bio-Rad μ -Radiance) using a $\times 20$ Nikon water/oil immersion objective. Sections with positive BSI lectin and SM α -actin staining and void of tissue folds from each region of the muscle were imaged and montaged using CorelDRAW 9 (Fig. 1). Each image montage was imported into Scion Image, and the total number of lectin-positive capillary vessels (determined by excluding any vessel with a lectin-labeled lumen diameter $> 10 \mu$ m), the total number of SM α -actin vessels (primarily arterioles and venules), the total number of muscle fibers, and total muscle cross-sectional area were measured for each cross section. The observer making the counts was blinded to the identity of the specimen. For soleus muscle end sections processed for myosin ATPase activity, entire cross sections were imaged with an Olympus BX51 microscope equipped with a DP70 digital camera using a $\times 4$ dry objective. To evaluate the percentage of type I slow oxidative muscle fibers, the number of dark muscle fibers was divided by the total number of muscle fibers.

Statistical analysis. Two-tailed Student's *t*-tests were used to compare histological parameters between control and mechanically stimulated groups per muscle region. Spatial differences in the microvascular response to mechanical vibration were based on assessing the interaction between region (end vs. mid) and the level of mechanical stimulation (control vs. vibration) with a two-factor ANOVA. Statistical significance was accepted for $P < 0.05$. All statistical analyses were performed using SigmaStat, and values are presented as means and SD.

RESULTS

Six weeks of extremely low-level, whole body vibrations for 15 min/day stimulated vascular remodeling in the soleus muscle. More specifically, in the end region of the mouse soleus muscle, high-frequency, low-magnitude vibrations superimposed on normal cage activity reduced the number of lectin-positive vessels per muscle fiber by 29% ($P = 0.02$; Fig. 2A) and SM α -actin-positive vessels per muscle fiber by 36% ($P = 0.05$; Fig. 3B) in the end region of the mouse soleus muscle.




Fig. 1. A: representative montage of a mouse soleus muscle cross section stained for BSI-lectin (green) and smooth muscle (SM) α -actin (red). Scale bar = 200 μ m. B: high-magnification image of soleus muscle stained for BSI-lectin (green) and SM α -actin (red). Scale bar = 20 μ m.

Although a similar trend was observed in the mid region of the soleus muscle, the vascular remodeling was not significant (Fig. 2, A and B). A two-way ANOVA revealed that, after allowing for effects of differences in region across end and mid regions, whole body mechanical vibration significantly reduced the number of lectin-positive vessels ($P = 0.012$) and SM α -actin-positive vessels ($P = 0.04$) in the soleus muscle. Mechanical vibrations did not alter muscle fibers per area or total area in either the end or mid regions of the muscle (Fig. 2, C and D). Quantification of the percentage of type I slow oxidative fibers in the end region of the mouse soleus muscle revealed that the whole body vibration stimulus did not induce slow-to-fast muscle fiber type transformation (Fig. 3).

DISCUSSION

The primary finding from this study is that brief exposure (15 min/day) of high-frequency (45 Hz), low-magnitude (0.3 g) mechanical vibration stimulates a reduction in the number of blood vessels per muscle fiber in mouse soleus muscle. Low level mechanical loading has been previously described to alter bone morphology and promote cellular growth in various

animals, including mice (17, 31, 33), and we report for the first time that this extremely small stimulus applied for as little as 15 min/day can produce microvascular alterations in mouse skeletal muscle.


Low-level vibrations decreased the number of lectin-positive vessels per muscle fiber in the end region of the muscle, yet did not influence muscle fibers per area or total cross-sectional area, thus indicating that these mechanical signals reduced the number of capillaries for a given muscle group. Again in the end region of the soleus muscle, the vibration decreased the ratio of SM α -actin-positive vessels per muscle fiber, which can be explained by the loss of lectin positive capillaries lined with SM α -actin-positive pericytes, the loss of existing arterioles and venules, or the dedifferentiation or loss of SM α -actin-positive perivascular cells along existing vessels. The microvascular adaptations, which could indicate a reduction in blood supply to the musculature responsible for posture, is an apparent paradox, given that this loading regimen has been shown to be anabolic in a range of species, including mice (17, 31, 33). The concept that the adaptive response of the muscle was more pronounced in the distal segment as contrasted to the midsection, although surprising, is consistent with evidence of stretch preferentially inducing new sarcomere growth near the muscle fiber end regions closest to the muscle-tendon joints and that the combination of stretch with electrical stimulation of the same fast contracting muscle causes slow-to-fast fiber type remodeling in the end region compared with the mid region (41).

High-frequency, low-magnitude vibrations in the same range as reported here can increase osteoblastic activity in adult animals, implicating this type of loading stimulation as a therapy for osteoporosis (29, 30, 33). In postmenopausal women and children with cerebral palsy, exposure of the skeleton to similar vibrations attenuates bone loss and increases bone mineral density (29, 34, 39). In this same group of mice, these mechanical signals served to enhance the trabecular bone morphology in the tibial metaphysis and epiphysis (42). Although the mechanisms by which bone cells sense such low-magnitude stimuli are not well understood, the data presented here indicate that the vasculature in a tissue functionally coupled to bone displays a similar sensitivity to this mechanical stimulus. The novelty of our result is that whole body vibration has the capability to affect the microvasculature and not solely that the number of vessels per muscle fiber is reduced in skeletal muscle for this particular application. Although other magnitudes, frequencies, and durations of vibration may cause different remodeling responses, this approach has indicated a mechanical means of modulating the microvasculature in muscle and may provide insight into the nature of the musculoskeletal "system." The significance of the current finding is that the highly noninvasive means of external whole body vibration has the potential to be used therapeutically or to help our understanding of vascular adaptations to mechanical loading regimens.

The direction of the reported microvasculature adaptations is surprising, given that this loading regimen has been shown to be anabolic to bone in a range of species, including mice (17, 31, 33). The reported reduction in the number of vessels per muscle fiber contradicts our original hypothesis. However, the directional effect of a mechanical stimulus depends on the resulting local stress alterations. For example, an increase in

AQ: 6

Fig. 2. Region-specific effects of whole body high-frequency, low-magnitude vibrations on soleus muscle: vibration effect on lectin-positive vessels per muscle fiber (A), vibration effect on SM α -actin-positive vessels per muscle fiber (B), vibration effect on muscle fibers per area (C), vibration effect on total cross sectional area (D). *Significant difference ($P < 0.05$) from age-matched control (AC) group using a Student's t -test [$n = 7$ for AC group, $n = 6$ for mechanically stimulated (MS) group]. Values are means and SD.



pressure in the case of hypertension might be thought to cause a positive vascular effect, but the pressure increase in smaller arterioles is accompanied by lumen reduction and an increase in wall thickness resulting in a reduction in wall stress. This local hemodynamic alteration is associated with microvascular rarefaction (23), demonstrating the need to understand the transduction of high-frequency, low-magnitude vibrations to the local stress level in blood vessels.


The cause and functional consequence of a vibration-induced reduction in the number of vessels per muscle fiber on blood supply or other physiological outcomes are unclear and could additionally be attributed to muscle fiber type remodeling or vibration-induced changes in the local extracellular environment. Previous work has linked high-frequency contractile stimuli with slow-to-fast fiber type remodeling (11, 20, 21, 40), suggesting an oxidative-to-glycolytic transformation. The present study suggests that low-level vibration loading does not affect muscle fiber number per area, muscle fiber size, or type I-to-type II transformation. However, the decrease in vessels per muscle fiber could still be, in part, explained by a transformation of fast oxidative to fast glycolytic fiber types. The resulting glycolytic fibers would require only a reduced number of capillaries, a finding consistent with the current data. A single cycle of tendon vibration for 15 min at 150 Hz

(1 mm displacement) of rat soleus muscle to decrease the secretion of bioassayable growth hormone secretion (12). Hudlicka et al. (15) suggest that electrical stimulation of afferent fibers in skeletal muscle leads directly to release of humoral factors, which then affect capillary growth. Finally, electrical stimulation producing contraction induced hyperfusion caused a threefold increase in VEGF mRNA abundance compared with passive hyperfusion (28). Thus the high-frequency, low-magnitude mechanical signals examined in this study could directly alter the local biochemical environment surrounding individual microvessels in the soleus muscle, which consists predominantly of oxidative fibers, causing a reduction in the number of vessels per muscle fiber.

Recent evidence showed that low-level mechanical signals can stimulate an anabolic response in bone (16, 30, 31, 33). The ability of whole body and tendon vibration to stimulate skeletal muscle to adapt (3, 12, 34, 38) further supports that muscle fibers are also sensitive to these vibratory signals. This report demonstrates the ability of extremely low-level vibrations applied for as little as 15 min/day to affect the microvascular architecture within skeletal muscle, but the functional outcome of these adaptations remains unclear. Our findings highlight the sensitivity of the microvasculature to subtle mechanical stimulation and the need to understand the potential system

ROF06

AQ: 7 Fig. 3. Effects of whole body, high-frequency, low-magnitude vibrations on end region muscle fiber type remodeling in mouse soleus muscle. A: percentage of type I muscle fibers in end region muscle sections indicated by a 4.3 pH myosin ATPase assay. B: representative image of a soleus end region cross section stained for myosin ATPase activity at 4.3 pH and counterstained with hematoxylin and eosin. Scale bar = 100 μ m.



level effects of whole body vibration. The current work suggests the possibility of developing noninvasive methods for manipulating the microvascular architecture in skeletal muscle.

GRANTS

This work was supported by National Heart, Lung, and Blood Institute Grants HL-65958 and HL-52309 to T. C. Skalak; US Army DAMD17-03-1-0777, Whitaker Foundation, RG-02-0564 to S. Judex; and National Institutes of Health AR-43498 to C. Rubin.

REFERENCES

1. **Badr I, Brown MD, Egginton S, Hudlicka O, Milkiewicz M, and Verhaeg J.** Differences in local environment determine the site of physiological angiogenesis in rat skeletal muscle. *Exp Physiol* 88: 565–568, 2003.
2. **Brown MD and Hudlicka O.** Modulation of physiological angiogenesis in skeletal muscle by mechanical forces: involvement of VEGF and metalloproteinases. *Angiogenesis* 6: 1–14, 2003.
3. **Burke D, Hagbarth KE, Lofstedt L, and Wallin BG.** The responses of human muscle spindle endings to vibration of non-contracting muscles. *J Physiol* 261: 673–693, 1976.
4. **Burke D, Hagbarth KE, Lofstedt L, and Wallin BG.** The responses of human muscle spindle endings to vibration during isometric contraction. *J Physiol* 261: 695–711, 1976.
5. **Clark CB, McKnight NL, and Frangos JA.** Strain and strain rate activation of G proteins in human endothelial cells. *Biochem Biophys Res Commun* 299: 258–262, 2002.
6. **De Ruiter CJ, Van Raak SM, Schilperoort JV, Hollander AP, and de Haan A.** The effects of 11 wk whole body vibration training on jump height, contractile properties and activation of human knee extensors. *Eur J Appl Physiol* 90: 595–600, 2003.
7. **Egginton S, Hudlicka O, Brown MD, Walter H, Weiss JB, and Bate A.** Capillary growth in relation to blood flow and performance in overloaded rat skeletal muscle. *J Appl Physiol* 85: 2025–2032, 1998.
8. **Falempin M and In-Albon SF.** Influence of brief daily tendon vibration on rat soleus muscle in non-weight-bearing situation. *J Appl Physiol* 87: 3–9, 1999.
9. **Fritton JC, Rubin CT, Qin YX, and McLeod KJ.** Whole-vibration in the skeleton: development of a resonance-based testing device. *Ann Biomed Eng* 25: 831–839, 1997.
10. **Fritton SP, McLeod KJ, and Rubin CT.** Quantifying the strain history of bone: spatial uniformity and self-similarity of low-magnitude strains. *J Biomech* 33: 317–325, 2000.
11. **Gorza L, Gundersen K, Lomo T, Schiaffino S, and Westgaard RH.** Slow-to-fast transformation of denervated soleus muscles by chronic high-frequency stimulation in the rat. *J Physiol* 402: 627–649, 1988.
12. **Gosselink KL, Roy RR, Zhong H, Grindeland RE, Bigbee AJ, and Edgerton VR.** Vibration-induced activation of muscle afferents modulates bioassayable growth hormone release. *J Appl Physiol* 96: 2097–2102, 2004.
13. **Hansen-Smith F, Egginton S, Zhou AL, and Hudlicka O.** Growth of arterioles precedes that of capillaries in stretch-induced angiogenesis in skeletal muscle. *Microvasc Res* 62: 1–14, 2001.
14. **Hansen-Smith FM, Watson L, Lu DY, and Goldstein I.** Griffonia simplicifolia I: fluorescent tracer for microcirculatory vessels in nonperfused thin muscles and sectioned muscle. *Microvasc Res* 36: 199–215, 1988.
15. **Hudlicka O, Graciotti L, Fulgenzi G, Brown MD, Egginton S, Milkiewicz M, and Granata AL.** The effect of chronic skeletal muscle stimulation on capillary growth in the rat: are sensory nerve fibres involved? *J Physiol* 546: 813–822, 2003.
16. **Judex S, Boyd SK, Qin YX, Turner S, Ye K, Müller R, and Rubin CT.** Adaptations of trabecular bone to low magnitude vibrations result in more uniform stress and strain under load. *Ann Biomed Eng* 31: 12–20, 2003.
17. **Judex S, Donahue LR, and Rubin C.** Genetic predisposition to low bone mass is paralleled by an enhanced sensitivity to signals anabolic to the skeleton. *FASEB J* 16: 1280–1282, 2002.
18. **Judex S, Garman R, Squire M, Busa B, Donahue LR, and Rubin C.** Genetically linked site-specificity of disuse osteoporosis. *J Bone Miner Res* 19: 607–613, 2004.
19. **Kakisis JD, Liapis CD, and Sumpio BE.** Effects of cyclic strain on vascular cells. *Endothelium* 11: 17–28, 2004.
20. **Lomo T, Westgaard RH, and Dahl HA.** Contractile properties of muscle: control by pattern of muscle activity in the rat. *Proc R Soc Lond B Biol Sci* 187: 99–103, 1974.
21. **Lomo T, Westgaard RH, and Engelbresten L.** Different stimulation patterns affect contractile properties of denervated rat soleus muscle. In: *The Plasticity of Muscle*, edited by D. Pette. New York: Walter de Gruyter, 1980.
22. **Matthews PB and Watson JD.** Action of vibration on the response of cat muscle spindle Ia afferents to low frequency sinusoidal stretching. *J Physiol* 317: 365–381, 1981.
23. **Prewitt RL, Rice DC, and Dobrian AD.** Adaptation of resistance arteries to increases in pressure. *Microcirculation* 9: 295–304, 2002.
24. **Price RJ and Skalak TC.** Chronic α 1-adrenergic blockade stimulates terminal and arcade arteriolar development. *Am J Physiol Heart Circ Physiol* 271: H752–H759, 1996.
25. **Pries AR and Secomb TW.** Structural adaptation of microvascular networks and development of hypertension. *Microcirculation* 9: 305–314, 2002.
26. **Prior BM, Lloyd PG, Yang HT, and Terjung RL.** Exercise-induced vascular remodeling. *Exerc Sport Sci Rev* 31: 26–33, 2003.
27. **Priplata AA, Niemi JB, Harry JD, Lipsitz LA, and Collins JJ.** Vibrating insoles and balance control in elderly people. *Lancet* 362: 1123–1124, 2003.
28. **Roca J, Gavin TP, Jordan M, Siafakas N, Wagner H, Benoit H, Breen E, and Wagner PD.** Angiogenic growth factor mRNA responses to passive and contraction-induced hyperperfusion in skeletal muscle. *J Appl Physiol* 85: 1142–1149, 1998.
29. **Rubin C, Recker R, Cullen D, Ryaby J, McCabe J, and McLeod K.** Prevention of postmenopausal bone loss by low-magnitude, high-frequency mechanical stimuli: a clinical trial assessing compliance, efficacy, and safety. *J Bone Miner Res* 19: 343–351, 2004.
30. **Rubin C, Turner AS, Bain S, Mallinckrodt C, McLeod and Anabolism K.** Low mechanical signals strengthen long bones. *Nature* 412: 603–604, 2001.
31. **Rubin C, Turner AS, Muller R, Mittra E, McLeod K, Lin W, and Qin YX.** Quantity and quality of trabecular bone in the femur are enhanced by a strongly anabolic, noninvasive mechanical intervention. *J Bone Miner Res* 17: 349–357, 2002.
32. **Rubin C, Xu G, and Judex S.** The anabolic activity of bone tissue, suppressed by disuse, is normalized by brief exposure to extremely low-magnitude mechanical stimuli. *FASEB J* 15: 2225–2229, 2001.
33. **Rubin CT, Sommerfeldt DW, Judex S, and Qin Y.** Inhibition of osteopenia by low magnitude, high-frequency mechanical stimuli. *Drug Discov Today* 6: 848–858, 2001.
34. **Russo CR, Lauretani F, Bandinelli S, Bartali B, Cavazzini C, Guralnik JM, and Ferrucci L.** High-frequency vibration training increases muscle power in postmenopausal women. *Arch Phys Med Rehabil* 84: 1854–1857, 2003.
35. **Skalak TC and Price RJ.** The role of mechanical stresses in microvascular remodeling. *Microcirculation* 3: 143–165, 1996.
36. **Torvinen S, Kannus P, Sievanen H, Jarvinen TA, Pasanen M, Kontulainen S, Nenonen A, Jarvinen TL, Paakkala T, Jarvinen M, and Vuori I.** Effect of 8-month vertical whole body vibration on bone, muscle performance, and body balance: a randomized controlled study. *J Bone Miner Res* 18: 876–885, 2003.
37. **Van Gieson EJ, Murfee WL, Skalak TC, and Price RJ.** Enhanced smooth muscle cell coverage of microvessels exposed to increased hemodynamic stresses in vivo. *Circ Res* 92: 929–936, 2003.
38. **Verschueren SM, Roelants M, Delecluse C, Swinnen S, Vanderschuere D, and Boonen S.** Effect of 6-month whole body vibration training on hip density, muscle strength, and postural control in postmenopausal women: a randomized controlled pilot study. *J Bone Miner Res* 19: 352–359, 2004.
39. **Ward K, Aslop C, Caulton J, Rubin C, Adams J, and Mughal Z.** Low magnitude mechanical loading is osteogenic in children with disabling conditions. *J Bone Miner Res* 19: 360–369, 2004.
40. **Westgaard RH and Lomo T.** Control of contractile properties within adaptive ranges by patterns of impulse activity in the rat. *J Neurosci* 8: 4415–4426, 1988.
41. **Williams P, Watt P, Bicik V, and Goldspink G.** Effect of stretch combined with electrical stimulation on the type of sarcomeres produced at the ends of muscle fibers. *Exp Neurol* 93: 500–509, 1986.
42. **Xie LQ, Donahue LR, Rubin C, and Judex S.** Low-level mechanical vibrations enhance bone morphology in the growing skeleton. In: *Proceedings of the Annual Fall Meeting of the Biomedical Engineering Society*, 2003.

AUTHOR QUERIES

AUTHOR PLEASE ANSWER ALL QUERIES

1

AQ1— Please return corrected printout of PDF (INCLUDING QUERY SHEET) within 48 hours via express delivery to: JAP Page Proof, 9650 Rockville Pike, Bethesda, MD 20814.

AQ8— You may delete e-mail address if you do not want it included.

AQ2— Please write out BSI.

AQ3— Key words duplicated from either title were deleted per journal style.

AQ4— APS Journal policy requires that a statement of protocol approval from an Institutional Review Board (IRB) or Institutional Animal Care and Use Committee (IACUC) or equivalent is to be included in the Methods section of all papers involving animal or human research. (See http://www.the-aps.org/publications/i4a/prep_manuscript.htm#humans_animals). Please provide us with the appropriate statement about institutional review of your study (IRB, IACUC, or equivalent) and indicate where specifically to add it on your proofs. Thank you.

AQ5— Are 9 and 17 reference nos. in sentence beginning “For mice in the MS group”?

AQ6— Sentence beginning “The significance of the current finding” edited for clarity; meaning intact?

AQ7— H&E written out okay in Fig. 3 legend?
